

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/68, G01N 33/53, 33/567, C12N 5/00	A1	(11) International Publication Number: WO 97/10361 (43) International Publication Date: 20 March 1997 (20.03.97)
(21) International Application Number: PCT/US96/14239 (22) International Filing Date: 4 September 1996 (04.09.96) (30) Priority Data: 9518611.0 12 September 1995 (12.09.95) GB 08/541,905 10 October 1995 (10.10.95) US (60) Parent Application or Grant (63) Related by Continuation US 08/541,905 (CIP) Filed on 10 October 1995 (10.10.95) (71) Applicant (for all designated States except US): LUDWIG INSTITUTE FOR CANCER RESEARCH [CH/US]; 1345 Avenue of the Americas, New York, NY 10105 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): VAN DAMME, Jo [BE/BE]; Tintoretterstraat 32, B-1040 Brussels (BE). PROOST, Paul [BE/BE]; Van Den Abeelelaan 21, B-3001 Heverlee (BE). HOUSSIAU, Frederic [BE/BE]; Avenue Ferdauci 36, B-1020 Brussels (BE). RENAULD, Jean- Christophe [BE/BE]; Boulevard General Wahis 32/28, B-1030 Brussels (BE).		(74) Agent: HANSON, Norman, D.; Felfe & Lynch, 805 Third Avenue, New York, NY 10022 (US). (81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: METHOD FOR MODULATING CELL APOPTOSIS (57) Abstract The invention is based upon the newly recognized ability of β chemokines to inhibit cell apoptosis. In particular, apoptosis of T cells is described. The known β chemokines I-309 and TCA-3 are examples of the β chemokines which inhibit apoptosis. One aspect of the invention is the use of these molecules to inhibit apoptosis. A second aspect of the invention is the use of β chemokine inhibitors or antagonists to provoke apoptosis.		

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

METHOD FOR MODULATING CELL APOPTOSIS**RELATED APPLICATIONS**

This is a continuation in part of Serial No. 08/541,905, filed October 10, 1995 and incorporated by reference.

FIELD OF THE INVENTION

5 The invention relates to the use of anti-apoptotic factors especially a mammalian anti-apoptotic factor which is a chemokine. More particularly the invention concerns the use of β chemokines such as I-309, and variants thereof, as
10 anti-apoptotic factors.

BACKGROUND OF THE INVENTION

Chemokines are a subset of cytokines. Cytokines are regulatory proteins secreted by white blood cells and by a variety of other cells in the body. The pleiotropic
15 actions of cytokines include numerous effects on cells in the immune system and the modulation of inflammatory responses ("The Cytokine Handbook", 2nd edition, 1994, ed. A. Thompson, Academic Press, p. 4). Chemokines are a
superfamily of proinflammatory cytokines. The chemokine
20 family consists of two branches. In the α chemokines, the first two cysteines are separated by a single amino acid residue (C-X-C) (e.g., IL-8 and platelet factor-4) whereas, in β chemokines such as TCA-3, the first two cysteines are
adjacent (C-C) (Lanig et al., 1994, *J. Immunol.* 153: 4625-
25 4635, incorporated by reference). Molecules in this family share at least 25% amino acid homology, have similar structures, and bind to seven-transmembrane-spanning
receptors of the rhodopsin superfamily. The genes encoding the α chemokines are located on human chromosome 4 and the
30 genes encoding the β chemokines are located on human chromosome 17.

Of the β chemokines, the murine protein TCA-3 was originally described by Burd et al. (1987, *J. Immunol.* 139:

3126-3131), which is incorporated by reference in its entirety. Its cDNA was cloned from a T cell helper clone library using a strategy aimed at identifying genes whose expression is induced upon stimulation. Two years later, Brown et al. (1989, *J. Immunol.* **142**: 679-687), incorporated by reference, similarly identified a cDNA called P500, which is identical to TCA-3, except for a 99 nucleotide insertion that was generated by alternative splicing in 8% of the cDNA clones that they analyzed. In this paper, TCA-3/P500 was also referred to as SISE (Small Inducible Secreted protein). The sequence of the human I-309 was reported by Miller et al. (1989, *J. Immunol.* **143**: 2907-2916), which is incorporated by reference in its entirety. Its cDNA was isolated from an IL-2 dependent T cell line using a similar strategy as reported in the preceding papers for TCA-3 and P500. The sequence of a genomic clone was reported later by Miller et al. (1990, *J. Immunol.* **145**: 2737-2744), incorporated by reference in its entirety, and the gene was shown to map to human chromosome 17 as expected for a β chemokine. The genomic structure supports the view that TCA-3 is the murine homologue of I-309.

There are three published reports describing the activity of an example of a β chemokine, that is human I-309, and the murine homologue TCA-3. These are cited herein, and are incorporated by reference. In the first report Wilson et al. (1990, *J. Immunol.* **145**: 2745-2750) have produced recombinant murine TCA-3 in CHO cells. They showed that intradermal injection of this material into mouse footpads results in a rapid swelling response, histologically characterized by a local accumulation of neutrophils.

In the second report Miller et al. (1992, *Proc. Natl. Acad. Sci. USA* **89**: 2950-2954) produced recombinant human I-309 in CHO cells and showed in vitro that purified I-309 stimulated migration of human monocytes but not neutrophils. I-309 was also shown to induce a transient

increase in cytoplasmic free calcium in monocytes but not in neutrophils or lymphocytes. The chemotactic activity required a 10 nM concentration (approximately 150 ng/ml) to be detectable. Maximal effect was found at 100-300 nM and half-maximal activity at approximately 20 nM.

Finally, in the third report Laning et al. (1994, *J. Immunol.* 153: 4625-4635) showed that a mouse tumor expressing TCA-3 lost its tumorigenicity, probably as a result of the anti-tumor immune response following the attraction of neutrophils and macrophages to the tumor site.

As previously mentioned, the human β chemokine I-309 was first described in 1989 by Miller et al. (1989, *J. Immunol.* 143: 2907-2916). The amino acid and cDNA sequences are described therein, along with other structural characteristics, such as a small polypeptide chain including a hydrophobic leader sequence which was thought to be involved in secretion of the corresponding protein. The function of the protein was unknown at that time, but it was suggested that because of the nature of the protein, that is, its relatively small size, secretion by T cells and other certain structural characteristics, it was a cytokine of unknown function. It has now been found that certain β cytokines, including P500 and TCA-3, I-309 demonstrate anti-apoptotic activity, when compared with the known anti-apoptotic agent IL-9, taught as a strong anti-apoptotic factor for thymic lymphoma cell lines. When one considers the effective chemotactic concentration of those proteins discussed infra, one concludes they are strong anti-apoptotic agents. Moreover, given the structural similarities between members of the β chemokine group and also the functional similarities, that is to say distinct chemokines have been shown to bind to the same receptor, it is considered that other members of the β chemokine group will also display anti-apoptotic activity. The following

references provide details of the β chemokine group and thus highlight the similarity between members of same. T. J. Schall (1991, *Cytokine* 3: 165-183); and Schall (1994, "The Cytokine Handbook", 2nd edition, ed. A. Thompson, Academic Press, pg. 419-460), incorporated by reference.

"Apoptosis", as used herein, describes one type of cell death. More particularly, it describes programmed cell death where, in the normal course of development, cells die and dismantle in a predictable and orderly fashion. For example, such cells tend to shrink within their membrane and their DNA is broken into nucleosomes for safe removal by cells of the immune system and in particular white blood cells.

The scientific community is only now recognizing the clinical importance of apoptosis and therefore only now beginning to invest in research that provides a greater understanding of the process and how the process can be advantageously manipulated in order to either promote, or in some instances, prevent cell death.

There are many examples of disorders and pathologies wherein apoptosis is implicated.

Heart attack is typically characterized by cardiac cell death and it therefore follows that factors that mitigate or prevent this cell death could be used to advantage to protect against the consequences of this condition. In addition, such factors could also be used to advantage in chemotherapy-induced cardiac tissue damage, or in stroke-induced tissue damage or even in kidney failure. The use of such factors would be particularly advantageous where an individual has a history of heart attacks or strokes or even belongs to a family having such a history.

Chemotherapy and radiotherapy also induce other forms of tissue damage and therefore it follows that co-administration or sequential administration of chemotherapy or radiotherapy with an anti-apoptotic factor may be used to advantage in order to mitigate or prevent this widespread damage.

In addition, premature and widespread apoptosis has been implicated in much of the damage associated with acquired immune deficiency syndrome (AIDS). It is thought that apoptosis may be a principal cause of death in uninfected T cells in AIDS patients, leading ultimately to the suppression of the patient's immune system. Thus factors which mitigate or prevent this form of cell death may have a role to play in preventing the damage that the AIDS virus causes in uninfected cells.

Cell death also occurs in instances of ischemia and thus the factors of the invention may also have a role to play in mitigating or preventing apoptosis in ischaemic organs.

In addition to uses of the invention to mitigate or prevent apoptotic activity, it is also apparent that one of the ramifications of the invention is the ability to promote apoptotic activity when desired. For example, the realization that chemokines, and in particular β chemokines such as I-309 and homologues in other species, have a role to play in preventing apoptosis, can be used to advantage in the manufacture of agents that selectively bind to this factor and thus suppress its anti-apoptotic function, thereby promoting cell death. Naturally occurring, artificial or synthetic agents may be used. For example, antibodies, monoclonal or polyclonal, can be made against the factor of the invention and then used to bind to same with a view to suppressing or annihilating its anti-apoptotic activity. Thus agents which selectively bind with the anti-apoptotic factor of the invention can be used

to promote apoptosis and moreover, having regard to the targeted delivery of such agents, can be used selectively to promote apoptosis of diseased tissue.

5 It may be advantageous to promote apoptosis in order to suppress or reverse tumor development. In this respect, it is of note that anti-apoptotic factors, for example, the product of the bcl-2 gene which encodes a cytoplasmic protein that protects against apoptosis without inducing proliferation by itself, has been shown to be involved in
10 some types of tumours.

In addition to the above, the invention can also be used in drug screening assays. For example, identifying chemokines, especially β chemokines in different parts of the body, either directly or indirectly, can be used to map
15 apoptosis in various parts of the body and thus areas, regions or organs most likely to suffer from cell death can be identified for drug screening purposes. In addition, the efficacy of apoptotic promoting agents and also agents acting as β chemokines, in particular I-309 antagonists,
20 can more accurately be determined by observing the activity of such agents in the presence of the factor of the invention. In addition, the invention can also be used to promote cell growth and proliferation and therefore can be used to provide proliferative tissue for drug screening
25 assays.

The full scope of the invention will be understood more fully following review of the disclosure which follows:

BRIEF DESCRIPTION OF THE FIGURES

30 Figure 1A shows anti apoptotic activity of various cell supernatants.

Figure 1B shows IL-6 content of the supernatants of figure 1A.

Figure 2 shows inhibition by anti IL-6 antibody of the activity of some, but not all fractions of activated CRL8066-MoT supernatants.

5 Figure 3 shows distinct fractionation of IL-6 and anti-apoptotic activity on phenyl sepharose.

Figure 4A depicts final purification of the anti-apoptotic factor on a Resource S column.

Figure 4B presents results of activity assay of various column fractions.

10 Figure 4C is an SDS-gel of various fractions.

Figure 5A shows anti apoptotic and chemotactic activities of purified I-309 on BW5147 cells, when dexamethasone is present, while figure 5B shows its activity in human THP1 monocytes.

15 Figure 6A sets forth data which shows that BW5147 cells proliferated when induced by recombinant I-309 expressed in COS cells, while figure 6B shows proliferation when stimulated by recombinant TCA 3 in the presence of dexamethasone.

20 Figure 7A sets forth evidence of recombinant I-309's anti-apoptotic activity on BW5147 cells.

Figure 7B presents electrophoretic data from DNA extracted from BW5147 cells, cultured in DEX with or without recombinant I-309.

25 Figures 8A and 8B present data showing inhibition of I-309 induced proliferation, in the presence of Bortadella pertussis toxin.

Figure 9 shows the anti-apoptotic activity of I-309 on thymic lymphoma cell lines 9T4A2 and NM3T2, following exposure to dexamethasone.

5 Figure 10 compares the anti-apoptotic effect of various chemokines.

Detailed Description of Preferred Embodiments

The invention concerns the anti-apoptotic activity of chemokines and in the following examples specific reference is made to the β chemokine I-309. Briefly, the
10 investigations described herein show that murine thymic lymphomas can be made to maintain significant proliferative activity and/or survival in the presence of the factor I-309. The anti-apoptotic activity of the factor is significant in that half maximal anti-apoptotic activity
15 was obtained as a concentration of only 1.6 ng/ml. This concentration is considerably lower (200 times lower) than the concentration required for monocyte chemotactic activity of this molecule suggesting that the primary function of this molecule may be suppression of cell
20 death.

In the following data specific reference is made to BW5147 murine thymic lymphoma. However it is of note that similar activity has also been obtained for two other lymphoma cell lines, i.e., 9T4A2 and NM3T2.

25 Materials and Methods

Cell cultures, cytokines and other reagents.

Thymic lymphoma cell line BW5147 (Ralph et al., 1973, *J. Immunol.* 110: 1470) was obtained from the American Type Culture Collection (ATCC). NM3T2 cell line was established
30 from a DBA-2 mouse treated with N-Methyl-N-nitrosourea (MNU) and 9T4A2 from a thymic lymphoma that spontaneously arose in an IL-9-transgenic mouse that had lost the IL-9

transgene (Renauld et al., 1994, *Oncogene* 9: 1327). These cells were cultured in Iscove's medium supplemented with 10% fetal calf serum, 1.5 mM L-glutamine, 0.24 mM L-asparagine, 0.55 mM L-arginine and 50 μ M 2-mercaptoethanol.

5 NK cell leukemia YT and T cell leukemia MoT/CRL8066 (Chen et al., 1983, *Nature* 305: 502-505) were obtained from the ATCC. Mo7E megakaryoblastic leukemia (Avanzi et al., 1988, *Br. J. Haematol.* 69: 359-366) was received from Dr. L. Pegoraro (University of Perugia). 5B12 and E5 are human T
10 cell clones derived from blood lymphocytes of healthy individuals and growing in the presence of PHA or allogenic feeder cells, PHA, IL-2 and IL-4.

Murine IL-9 was produced by expression in baculovirus and purified by affinity chromatography in our laboratory.
15 Saturating doses (200U/ml) were used in every experiment. Recombinant TCA-3 was purchased from R&D laboratories and Pertussis Toxin was from BIOMOL (Plymouth Meeting, PA). Dexamethasone (DEX) was obtained from Sigma and used at 0.25 μ M concentration in all experiments. The rat anti-
20 murine IL-6 receptor, 15A7 (Coulie et al, 1990, *Curr. Top. Microbiol. Immunol.* 166: 43-46), and mouse anti-human IL-6, AH64 (Brailly et al, 1994, *Clin. Chem.* 40: 116-123), antibodies were used at 50ug/ml and at 1/100 dilution of ascites fluid, respectively.

25 The 7TD1 bioassay used to measure IL-6 was performed as described previously (Van Snick et al., 1986, *Proc. Natl. Acad. Sci. USA* 83: 9679-9683). Briefly, 2×10^3 7TD1 cells were seeded in microwells and cultured for 4 days in the presence of serial dilutions of the supernatants. The
30 proliferation was evaluated by colorimetric determination of hexosamindase levels as described by Landegren et al., 1984, *Cytokine* 3: 165-183.

Anti-apoptotic bioassays

Cell viability was determined using a propidium iodide incorporation assay as described (Lee et al., 1993, *J. Immunol.* **151**: 5208). Briefly, cells were incubated for 30 minutes with propidium iodide (125 μ g/ml) at room temperature before FACS analysis with a FACScan flow cytometer (Becton-Dickinson). Under these conditions, dead cells are brightly stained while live cells are not. A minimum of 5000 cells were counted per sample.

To test the influence of cytokines on proliferation in the presence of DEX, BW5147 cells were incubated in triplicate microtiter wells in the presence of DEX at a final 0.25×10^{-6} molar concentration with or without cytokines, in a 200 μ l volume. Cultures were pulsed 2-4 days later for 6 h with 0.5 μ Ci 3 H-thymidine. Alternatively, cell proliferation was measured by an hexosaminidase assay, as described (Uyttenhove et al., 1988, *Proc. Natl. Acad. Sci. USA* **85**: 6934).

Purification procedures

MoT/CRL8066, a hairy cell leukemia T type, was cultured in RPMI-1640 medium supplemented with 10% heat-inactivated (56°C, 30 min) FCS, 0.55 mM L-arginine, 0.24 mM L-asparagine, and 1.25 mM L-glutamine. For the production of large batches, cells were washed, resuspended in the same medium containing 5% FCS and stimulated at a concentration of 5×10^5 cells/ml with PMA (50ng/ml) (Sigma). After heating for 30 minutes at 57°C in order to inactivate HTLV-II virus, supernatants were buffered with sodium phosphate, pH 7.0, and ammonium sulfate was added to a final concentration of 1.2 M. This preparation was chromatographed on a Phenyl-Sepharose Fast flow column (Pharmacia) at 8 ml/min. After a 500ml wash with 1.2 M ammonium sulfate, active material was recovered by a two-step elution at 3 ml/ml with 1 liter of 0.7 M and 0.5 M ammonium sulfate.

After concentration, the material was transferred into a sodium acetate 50 mM, pH 5.5 buffer containing 0.01% (vol/vol) Tween 20 before cation exchange chromatography on a Sulfo-Propyl Sepharose Fast flow gel (Pharmacia) at 3 ml/min. After washes with 100 ml of the same buffer and 140 ml of 0.3 M NaCl in the same buffer, elution was performed with a 300 ml linear gradient of NaCl to 1 M, the active material being eluted at 0.5 to 0.75 M NaCl.

The active fractions were concentrated, transferred into 50 mM phosphate buffer, pH 7.0, containing 1 M NaCl and 0.01% (vol/vol) Tween 20 and fractionated on a Superdex 70 size-exclusion column (Pharmacia). The active material (eluted with apparent molecular weight of 12-16 kD) was transferred into 100 mM Na₂HPO₄ buffer pH 7.0 containing 1 M Na₂SO₄ and loaded onto a TSK-phenyl column (LKB, Bromma, Sweden) at 0.5 ml/min. After a 5 min wash in the starting buffer, elution was carried out at 0.5 ml/min with a linear gradient of a 1/1 mixture of sodium phosphate (0.1 M, pH 7.0) and ethylene glycol (from 0 to 60% in 30 min), the active material being eluted at 30 % of this buffer.

The active fractions were transferred into sodium acetate 50 mM, pH 5.5 buffer containing 0.01 % (vol/vol) Tween 20 before cation exchange chromatography on a Sulfo-Propyl Resource S column (Pharmacia) at 0.8 ml/min. Elution was performed at 0.8 ml/min with a 10 min linear gradient of NaCl to 0.2 M, followed by a 30 min gradient to 0.4 M NaCl, the active material being eluted at 0.3 M NaCl.

The most active fraction was adjusted to contain 0.1 % trifluoroacetic acid and further analyzed by reverse phase chromatography on a Vydack C4 column. The column was developed at 0.8 ml/min with a 40 min linear gradient from 8 to 56 % acetonitrile in 0.1 % trifluoroacetic acid. Fractions were collected in Eppendorf tubes containing 5 µl of Tween 20 (1% in water), lyophilized and resuspended in

Na₂HPO₄ buffer (50 mM, pH 7.0) before testing the anti-apoptotic activity.

cDNA cloning and COS cell transfections

The coding sequence of the human I-309 cDNA was amplified
5 by RT-PCR using total RNA extracted from a PHA activated T
cell clone (E5). Reverse transcription was performed on 5
µg total RNA with an oligo(dT) primer, cDNA corresponding
to 200 ng of total RNA was amplified for 30 cycles by PCR
with the following specific primers into which an EcoRI
10 site was inserted: sense 5'-TCCAGGAATTCCCAAGCCAGACCAGAA-3'
(SEQ ID NO:1) (from position 19 of the cDNA sequence),
antisense 5'-TTGTAGAATTCAAATGTTTAAAGTGCAACA-3' (SEQ ID
NO:2) (from position 503 of the cDNA sequence). The
amplification product was digested by EcoRI and cloned in
15 the pCDSRα expression vector (Takebe et al, 1988) The
nucleotide sequence was determined by the dideoxynucleotide
chain termination method using the Taq cycle sequencing
system (Amersham) with ³²P-ATP labelled oligonucleotides,
and no difference was observed with the published I-309
20 sequence (Genbank Accession Number M57502) (Miller et al.,
1989, *J. Immunol.* **143**: 2907-2916).

Anti-apoptotic activity and IL-6 content of various human cell supernatants. (Figure 1)

A: For the anti-apoptotic assay, BW5147 cells (10³/well)
25 were incubated for 3 days in the presence of
dexamethasone (DEX) (0.25 µM) with or without murine
IL-9 (200 U/ml) or 3.3% of supernatant from the Mo7E
megakaryoblastic leukemia cell line, MoT/CRL8066 T
cell leukemia cell line (stimulated or not with PMA
and IL-2 for 3 days) the YT NK cell leukemia
30 stimulated with PMA and IL-2 or 10% of T cell clone
supernatants (5B12 and E5). Cell proliferation was
evaluated by measuring thymidine incorporation and
standard deviations were calculated from triplicate

cultures. In the absence of DEX, thymidine incorporation was $117,947 \pm 7,122$ cpm.

5 B: The IL-6 content of the supernatants was measured by the 7TD1 bioassay as described. 1 U/ml corresponds, to the concentration required for half-maximal proliferation (approximately 5 pg/ml for human IL-6).

Inhibition by anti-IL-6 antibody of the activity of some but not all fractions of activated CRL8066-MoT supernatants. (Figure 2)

10 BW5147 cells (10^3 /well) were incubated for 3 days in the presence of DEX ($0.25 \mu\text{M}$) with or without human IL-6 (500 U/ml) or aliquots from elution fractions of the Superose chromatography. The rat anti-murine IL-6 receptor, 15A7, and the mouse anti-human IL-6 antibodies were used at 50
15 $\mu\text{g/ml}$ and at 1/100 dilution of ascites fluid, respectively. Cell proliferation was evaluated by measuring thymidine incorporation and standard deviations were calculated from triplicate cultures.

Distinct fractionation of IL-6 and anti-apoptotic activity on phenyl sepharose. (Figure 3)

20 Supernatants from PMA-stimulated MoT/CRL8066 cells were heated for 30 minutes at 57°C in order to inactivate HTLV-II virus, buffered with sodium phosphate, pH 7.0 before ammonium sulfate was added to a final concentration of 1.2
25 M and loading on a Phenyl-Sepharose Fast flow column (Pharmacia). After appropriate washing, a three step elution was performed with 0.75, 0.5, and 0.0 M ammonium sulfate and the activity of the fractions was evaluated in the 7TD1 assay for IL-6 and in the BW5147 anti-apoptotic
30 assay as described in Material and Methods.

Final purification of anti-apoptotic factor on Resource S column and SDS PAGE gel. (Figure 4)

5 A: After three successive steps of separation on a phenyl-sepharose gel, Sulfo-Propyl column and Superdex size exclusion column, the active fractions were fractionated by cation-exchange chromatography on a Sulfo-Propyl Resource S column. Elution was performed by a NaCl gradient as shown by the dotted line (Buffer B = 1M NaCl). The optical density (280 nm) of the
10 eluted material (stripplled area) and the anti-apoptotic activity (closed squares) are shown. 1 U/ml of anti-apoptotic activity is arbitrarily defined as the concentration giving half-maximal protection against DEX in the BW5147 bioassay described in
15 Materials and Methods.

B: As fractions were removed from the Resource S cation exchange column, they were tested for anti-apoptotic activity in the anti-apoptotic assay described supra. The results, presented in figure 4B, show that elution
20 fraction 21 was, in fact, the fraction containing the apoptotic activity. Several of the fractions were subjected to standard SDS-PAGE analysis, and the results are shown in figure 4C.

25 *Anti-apoptotic and chemotactic activities of purified I-309. (Figure 5)*

A: BW5147 cells (10^3 /well) were incubated for 3 days in the presence of DEX ($0.25 \mu\text{M}$) with or without serial dilutions from fraction 21 of the Resource S chromatography shown in Figure 4. Cell proliferation
30 was estimated by a colorimetric hexosaminidase assay. I-309 concentrations were deduced from the micro-sequencing analysis.

B: The same method was used in a chemotactic assay using the human THP1 monocytic cell line (closed symbols).

Results are indicated as chemotactic indices. Purified human MCP-3 was used as a positive control for this assay (open symbols).

5 *Proliferation of BW5147 cells induced by recombinant I-309 expressed in COS cells and by recombinant TCA-3 in the presence of dexamethasone. (Figure 6)*

10 A: BW5147 cells (10^3 /well) were incubated for 3 days in the presence of DEX ($0.25 \mu\text{M}$) with or without a 1/15 dilution of supernatant from COS cells transfected with the I-309 cDNA or a control plasmid. Cell proliferation was evaluated by measuring thymidine incorporation and standard deviations were calculated from triplicate cultures.

15 B: The anti-apoptotic activity of serial dilutions of recombinant murine TCA-3 was measured as described in A.

20 Further work was then carried out using recombinant I-309 on BW5147 cells. In these experiments, the BW 5147 cells (again, 10^3 /well), were incubated for 3 days in the presence of 0.25 mM DEX, either with or without purified recombinant I-309 (50 ng/ml). By measuring thymidine incorporation, cell proliferation was evaluated. Triplicate cultures were used to develop the data and to calculate standard deviations. The data are presented in figure 7A. In figure 7B, the results of electrophoresis of DNA which was extracted from BW5147 cells which had been cultured for 24 hours in the presence of DEX with or without 400 ng/ml of recombinant I-309 are shown.

30 *Inhibition by Pertussis toxin of I-309-induced proliferation of BW5147 cells in the presence of dexamethasone. (Figure 8)*

BW5147 cells (10^3 /well) were incubated for 3 days in the presence of DEX ($0.25 \mu\text{M}$) with or without a 1/15 dilution of supernatant from COS cells transfected with the I-309 cDNA or a saturating concentration of mIL-9 (200U/ml).
5 Cell proliferation was evaluated by measuring thymidine incorporation and standard deviations were calculated from triplicate cultures.

In a similar experiment, DEX, at 0.25 mM , with or without 5% activated Mo cell supernatant and with or without B. pertussis toxin (2.5 ng/ml), or anti human IL-6 mAb AH65 (ascites fluid, 0.5%), were added. Thymidine incorporation is shown in figure 8B.
10

Anti-apoptotic activity of I-309 for 9T4A2 and NM3T2 thymic lymphoma cell lines exposed to dexamethasone. (Figure 9)
15 Cells were incubated with dexamethasone in the presence or absence of I-309 (100 U/ml) or IL-9 (500 U/ml) for 20 h. Cell viability was measured by FACS analysis after staining with propidium iodide as described in Material and Methods.

A series of experiments were carried out in which the antiapoptotic activity of various human chemokines on BW5147 was tested. In each case, BW5147 cells (10^3 /well) were incubated for 3 days in the presence of DEX(0.25 mM), either with or without I-309 (40 ng/ml), MCP-1(25 ng/ml), MCP-2 (250 ng/ml), MCP-3 (250 ng/ml), RANTES (250 ng/ml),
20 IL-8 (125 ng/ml), NAP-2 (125 ng/ml), granulocyte chemotactic protein -2 (250 ng/ml), and macrophage inflammatory protein 1a (125 ng/ml). Cell proliferation was measured in a thymidine incorporation assay, using triplicate cultures, all as described supra. Figure 10
25 presents these results.
30

Results

Identification of anti-apoptotic activity distinct from IL-6 in human T-cell supernatants

In a previous report, it had been shown that the murine
5 thymic lymphoma BW5147 is protected against dexamethasone-
induced apoptosis by IL-9, IL-4 and to a lesser extent by
IL-6. By contrast, IL-1, IL-2, IL-7, IL-10, IL-13, IFN γ
and TNF α were not active in this assay (Renauld *et al*,
1995, *Blood* **85**: 1300-1305), as well as IL-3, IL-11, LIF,
10 GM-CSF and Steel Factor (unpublished data). Since human
IL-9 and human IL-4 are not active on murine cells, we used
this bioassay to screen a series of human supernatants for
another anti-apoptotic activity. Since human IL-6 is
active on murine cells, we simultaneously measured the IL-6
15 titers of these supernatants. As shown in Figure 1A,
proliferation of BW5147 cells cultured in the presence of
dexamethasone was observed with murine IL-9 but also with
some human cell supernatants such as those from activated
T cell leukaemia (CRL8066-MoT) or from CD4+ T cell clones
20 (5B12, E5). Interestingly, this activity was not strictly
correlated with the concentration of IL-6 in these
supernatants as shown in Figure 1B. Although the most
potent supernatant (activated CRL8066-MoT) contained large
amounts of IL-6, other active samples (5B12 and E5
25 supernatants) were essentially devoid of IL-6, suggesting
that a factor distinct from the previously checked
cytokines could protect BW5147 cells against dexamethasone-
induced apoptosis.

To further discriminate this putative factor from IL-6,
30 CRL8066-MoT cell supernatants, concentrated by adsorption
to sulfopropyl sephadex beads, were chromatographed on a
size exclusion gel by FPLC. This experiment showed that a
factor smaller in size than IL-6 and not inhibited by anti-
human IL-6 or anti-mouse IL-6 receptor antibodies was also
35 active in this assay (Figure 2). This conclusion was
confirmed by results obtained in a hydrophobic interaction

chromatography on phenyl sepharose. As shown in Figure 3, the major anti-apoptotic activity was eluted at 0.75 M ammonium sulphate, while IL-6 was recovered only at 0.0 M salt.

5 Purification and micro-sequencing of the factor responsible
 for the anti-apoptotic activity

CRL8066-MoT cells were produced by stimulating the cells at a concentration of 500,000 cells/ml with PMA at 50 ng/ml for 3 days in medium containing 5% FCS. After
10 chromatography on a Phenyl-Sepharose column, the material was fractionated on a cation exchange gel at pH 5. The active fractions were transferred into a neutral phosphate buffer containing 1 M NaCl and a 10^{-4} dilution of Tween 20, and subjected to gel filtration chromatography. The anti-
15 apoptotic activity eluted with an apparent molecular weight ranging from 10 to 20 kD with a peak at 15 kD. Further fractionation was carried out by hydrophobic interaction chromatography on a TSK phenyl column from which the active material eluted at 0.7 M Na_2SO_4 and 15% ethylene glycol.
20 Finally, the active fractions were purified to homogeneity on a Resource S cation exchange column by FPLC. The results of this last chromatography step are shown in Figure 4, together with the corresponding SDS PAGE analysis of the active fractions.

25 The purity of this preparation was confirmed by the fact that HPLC analysis of the most active fraction (fraction 21, see Figure 4) showed a single protein peak, from which the anti-apoptotic activity was retrieved. Sequencing of 30 pmoles of protein from fraction 21 of the Resource S column
30 (Figure 4) generated the following 27 amino acid sequence:
NH₂-Lys-Ser-Met-Gln-Val-Pro-Phe-Ser-Arg-Cys-Cys-Phe-Ser-Phe-Ala-Glu-Gln-Glu-Ile-Pro-Leu-Arg-Ala-Ile-Leu-Cys-Tyr, (SEQ ID NO:3) which was completely identical to amino acids 1 to 27 of the β -chemokine I-309 (Miller et al, 1989, J.

Immunol. **143**: 2907-2916; Miller *et al.*, 1990, *J. Immunol* **145**: 2737-2744).

Comparison of the specific activity of I-309 in chemotactic and anti-apoptotic in vitro assays.

5 Based on concentrations deduced from the micro-sequencing analysis, the specific activity of purified I-309 was measured in the anti-apoptotic assay. Maximal protective activity for BW5147 cells was reached at a 3 ng/ml concentration (Figure 5A). If, in this assay, one defines
10 one unit as the concentration required to give half-maximal protection against dexamethasone, one unit would be equivalent to 1.65 ng/ml (approximately 100pM). Interestingly, this specific activity was found to be in a similar range as that of murine IL-4 in the same bioassay
15 (Renauld *et al.*, 1995, *Blood* **85**: 1300-1305).

The same purified material was tested in a chemotactic assay using the human monocytic cell line THP1. As shown in Figure 5B, significant chemotactic responses to I-309 required considerably higher concentrations. Similar
20 results were obtained using normal monocytes (data not shown). Taken together, these data indicate that the anti-apoptotic effect reported here is a much more sensitive assay and that I-309 might be in fact much more effective in vivo as an anti-apoptotic agent than as a chemotactic
25 factor.

Anti-apoptotic activity of recombinant I-309 and TCA-3.

To formally establish that the anti-apoptotic activity observed with our purified material can be ascribed to I-309 cDNA was amplified by RT-PCR from a PHA activated T
30 cell clone, inserted into an expression vector and transfected into COS-7 cells. As shown in Figure 6, the supernatant from I-309-transfected COS cells strongly stimulated proliferation of BW5147 cells in the presence of dexamethasone, while control COS cell supernatant was

totally inactive. In addition, baculovirus derived TCA-3, considered as the murine homologue of I-309, had a similar activity with half-maximal effect observed at approximately 0.5 ng/ml.

5 Inhibition of the anti-apoptotic activity of I-309 by
 Pertussis toxin.

The receptor molecule for I-309/TCA-3 has not yet been identified. However, every chemokine receptor identified so far consist in a transmembrane protein with seven
10 transmembrane domains that transduce the signal through G proteins. Since Pertussis toxin is known to interfere with the activation mediated by a subset of these receptors, we analyzed the effect of this molecule on the anti-apoptotic activity of I-309. As indicated in Figure 8, when BW5147
15 cells were exposed to dexamethasone in the presence of recombinant I-309, Pertussis toxin completely abrogated the proliferation induced by I-309 in the presence of DEX. By contrast, Pertussis toxin did not inhibit the IL-9-induced cell proliferation, demonstrating that completely distinct
20 mechanisms of signal transduction are involved. These results also suggest that chemokine receptors linked to G proteins may play an unexpected role in the control of cell survival under apoptosis-promoting conditions.

25 Anti-apoptotic activity of I-309 on other thymic lymphoma
 cell lines.

To see whether the anti-apoptotic activity of I-309/TCA-3 can be extended to other thymic lymphoma cell lines, we tested the activity of purified I-309 on two other dexamethasone-sensitive cell lines, 9T4A2 and NM3T2, by
30 measuring cell death 20 hours after exposure to dexamethasone. As shown in Figure 9, I-309 was found to be as potent as IL-9 in inhibiting apoptosis in these cells. These results suggest that the anti-apoptotic activity of I-309 may be a common phenomenon for these types of tumor
35 lines.

The foregoing examples and disclosure describe the invention, which is a method of modulating cell apoptosis. In one aspect of the invention, the modulation involves the inhibition of cell apoptosis by administering an amount of

5 a β chemokine to a cell sample, in an amount sufficient to inhibit apoptosis of cells in said sample. Especially preferred are proteins which have I-309 activity including mammalian analogous of I-309, such as TCA-3, P500, and so forth, as well as anti-apoptotic fragments of these β

10 chemokines. Especially preferred are methods of inhibiting T cell apoptosis, using the β chemokines in the manner described. "Effective amount" as used herein refers to any amount of the β chemokine sufficient to inhibit cell apoptosis. Preferably, the amount of β chemokine used will

15 range from about 0.1 ng/ml up to about 100 ng/ml. Most preferably, the amount dosed will range from about 1.0 ng/ml to about 3.0 ng/ml. When used in vivo, the protein may be administered in accordance with any of the standard therapeutic regimes, such as orally, intravenously,

20 intramuscularly, subcutaneously, intranasally, or intradermally, with or without a carrier, adjuvant, or additional material to protect the active ingredient from degradation by normal physiological processes. When used in vitro, solutions, emulsion, etc. may be used to

25 facilitate the applications of the material.

One may also inhibit cell apoptosis, as indicated herein, by administering antagonists of β chemokines, such as I-309 antagonists, to a subject. As indicated, these antagonists may be antibodies, such as polyclonal or monoclonal

30 antibodies, soluble forms of the I-309 receptor, and also derivatives of the β chemokines which share sufficient structural similarities with the β chemokines to bind to β chemokine receptors, but sufficiently different so as to be apoptotically inert.

Other aspects of the invention will be clear to the skilled artisan, and need not be repeated here.



5 The terms and expressions which have been employed are used as terms of description and not of limitation, and there is no intention in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof, it being recognized that various modifications are possible within the scope of the invention.

Claims:



1. Method for inhibiting apoptosis, comprising administering to a cell sample an amount of an apoptosis inhibiting β chemokine sufficient to inhibit apoptosis of cell in said cell sample.
2. The method of claim 1, wherein said cells are T cells.
3. The method of claim 1, wherein said β chemokine is a mammalian β chemokine.
4. The method of claim 3, wherein said mammalian β chemokine possesses I-309 activity.
5. The method of claim 4, wherein said β chemokine is I-309.
6. The method of claim 1, wherein said β chemokine is administered in an amount ranging from about 0.1 ng/ml to about 100 ng/ml.
7. The method of claim 6, wherein said β chemokine is administered in an amount ranging from about 1.0 ng/ml to about 3.0 ng/ml.
8. The method of claim 1, comprising administering said β chemokine to a subject in vivo.
9. The method of claim 1, comprising administering said β chemokine to a culture of cells in vitro.
10. Method for enhancing cell apoptosis comprising administering to a sample of cells an amount of an inhibitor of β chemokine sufficient to enhance apoptosis of cells in said sample.

11. The method of claim 10, wherein said inhibitor is an inhibitor of a β chemokine which has I-309 activity.
12. The method of claim 10, wherein said inhibitor is an antibody which specifically binds to said β chemokine.
13. The method of claim 10, wherein said inhibitor is a soluble form of a receptor for said β chemokine.

1/10


FIG. 1A**FIG. 1B**

2/10



FIG. 2**FIG. 3**

3/10



FIG. 4A





4/10

FIG. 4B**FIG. 4C**


5/10

FIG. 5A**FIG. 5B**

6/10



FIG. 6A**FIG. 6B**

7/10



FIG. 7A**FIG. 7B**

DEX	+	+
I-309	-	+

8/10


FIG. 8A**FIG. 8B**

9/10

FIG 9A**FIG 9B**

10/10

FIG. 10



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/14239

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12Q 1/68; G01N 33/53, 33/567; C12N 5/00

US CL : 435/6, 7.1, 7.21, 240.2,

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6, 7.1, 7.21, 240.2,

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, MEDLINE, CAPLUS, BIOSIS, EMBASE, SICSEARCH, WPIDS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X -- Y	LANING et al. Inhibition of In Vivo Tumor Growth by the B Chemokine, TCA3. J. Immunol. 1994, Vol. 153, pages 4625-4635. See entire document.	1-4, 8, 9, 11 -- 5-7, 10, 12, 13
X -- Y	LUO et al. Biologic Activities of the Murine B-Chemokine TCA3. J. Immunol. 1994, Vol. 153, pages 4616-4624. See entire document.	1-4, 6-12 -- 5, 13

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

11 OCTOBER 1996

Date of mailing of the international search report

28 OCT 1996

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JHUNG-WON COLBY, PH.D.

Telephone No. (703) 308-0196

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)